

The zooplankton from some lakes and ponds in North Albania with different size and altitude

Dimitar KOZUHAROV

The data about the zooplankton in the fresh waters of Albania are not many. The greater part of the investigations has been done on the Scutari lake because of its size - 370 km². and the transborder attitude. The investigations on the crustaceans of the Skutari lake particularly started in the beginning of the century (BREM & ZADERBAUER, 1905; VERESHCHAGIN, 1912; GESSNER, 1934; NEDELKOVIC, 1959; PETKOVSKI, 1961). There are no data about the zooplankton of the prevailing part of the numerous small lakes and ponds in the country. There are no data about the rotifers from the Skutari lake either. Eight species of Cyclopoids from the Albanian ground waters were reported by PANDOURSKI (1977).

In July 1995 during a speleological expedition of the Bulgarian Federation of Speleology, zooplankton samples from eight lakes and ponds different in size and hydrological regime, were collected (Fig. 1). The sampling was carried out by an „Apstein“ net with 40mkm measure. In the littoral zone of the lakes a hand net was used for the collection of littoral animals.

Descriptions of the investigated biotops and the methods used for taking samples:

1, 2 Samples - Scutari Lake - geographic position 42.4 N, 19.3E - Surface area 370km². 16 m a.s.l., SE coast, first sample taken by a plankton net, second sample taken by a hand net in the littoral zone. Sampling date - 23.06.1995

3 Sample - Artificial lake for agricultural irrigation in the Vrithi village 350 m.a.s.l. - 15.06.1995

4 Sample - Artificial lake for watering above Vrithi village. 1200 m.a.s.l. Sampling date - 15.06.1995

5 Sample - Artificial lake for watering below Vrithi village. Sampling date - 16.06.1995




Fig.1. Map of Albania and the investigated region - marked in the rectangle

Table 1

The invertebrate taxa found in the investigated lakes and ponds in the North Albania

Legend: 1, 2 - Scutari lake; 3 - Artificial lake for agricultural irrigation in the Vrithi village; 4 - Artificial lake for watering above Vrithi village; 5 - Artificial lake for watering below Vrithi village; 6, 7 - Artificial lake for watering at the Stani I Drugomiri place; 8, 9 - Source at the Proni e Shkurtit place; 10 - Little well at the Stani I Drugomiri place

Taxa	1	2	3	4	5	6	7	8	9	10
Rotatoria										
<i>Lecane (M.) arcuata</i> (Bryce, 1891)		+								
<i>Polyarthra remata</i> Skorikov, 1896	+		+							
<i>Polyarthra</i> sp.	+									
<i>Asplanchna sieboldi</i> (Leidig, 1854)			+							
<i>Asplanchna</i> sp.							+			
<i>Synchaeta</i> sp.							+	+		
<i>Brachionus urceus</i> (L., 1758)							+			
<i>B. urceus sericus</i> Rouselet, 1907							+			
<i>Keratella cochlearis</i> (Gosse, 1851)	+									
<i>K. tecta</i> (Gosse, 1851)	+									
<i>K. quadrata frenzeli</i> (Eckstein, 1895)	+									
<i>Euehlanis dilatata</i> Ehrenberg, 1832	+	+								
<i>Kellicottia longispina</i> (Kellicott, 1879)	+									
<i>Testudinella</i> sp.							+			
Cladocera										
<i>Sida cristalina</i> (O.F. Muller, 1776)	+	+								
<i>Daphnia similis</i> Claus, 1876							+			
<i>D. pulex pulex</i> (De Geer, 1835)			+		+	+				
<i>D. galeata</i> Richard, 1896						+				
<i>Daphnia</i> sp. juv.									+	
<i>D. gr. pulex</i> - ehipia			+		+	+				
<i>D. gr. longispina</i> - ehipia						+	+			
<i>Moina brachiata</i> (Jurine, 1820)			+		+					
<i>Bosmina longirostris</i> (O.F. Muller, 1776)	+		+	+						
<i>Acroperus harpae</i> (Baird, 1835)		+								
<i>Alona gutata</i> Sars, 1862		+								
<i>A. rectangula</i> Sars, 1862		+								
<i>Chydorus sphaericus</i> (Muller, 1785)		+								
Copepoda										
<i>Eudiaptomus</i> sp. juv.	+									
<i>Eucyclops serrulatus</i> (Fischer, 1853)		+		+						
<i>E. serrulatus proximus</i> (Fischer, 1853)						+	+			
<i>Eucyclops</i> sp.		+								
<i>Trophocyclops prasinus</i> (Fischer, 1860)			+							
<i>Cyclops vicinus</i> Uljanin, 1875	+									
<i>Acanthocyclops vernalis</i> (Fischer, 1860)				+	+	+				
Copepodites	+	+	+	+						

Taxa	1	2	3	4	5	6	7	8	9	10
Nauplii	+	+		+		+				
Others										
<i>Gammarus</i> sp.										+
Asellidae gen. sp.									+	
Coleoptera <i>Hydrous</i> sp.							+			
Fam. Chironomidae larvae	+	+							+	
<i>Salamandra</i> larvae									+	
Planaria gen. sp.								+		

6, 7 Samples - Artificial lake for watering, above Rasma village at the Stani I Drugomiri place - 1590 m. a.s.l. - two samples. First taken by a plankton net, second taken by a hand net. Sampling date - 20.06.1995

8, 9 Samples - Source at the Proni e Shkurtit place 1460 m. a.s.l. - two samples, first by plankton net, second by hand net. Sampling date - 21.06.1995.

10 Sample - Little well at the Stani I Drugomiri place. 1590 m.a.s.l. Sampling date - 23.06.1995

All the investigated artificial ponds have an area between 100 and 1000 m².

Of all of the investigated waters 14 taxa Rotatoria, 11 taxa Cladocera and 6 taxa Copepoda were established (Table 1). The Cladocera *Daphnia galeata* found in the artificial lake N 6 was not reported for Albania before. PETKOVSKI (1961) reported *Daphnia cuculata* kahlbergensis from the Skutari lake. This species belongs to the same species group without seta comb of longer setae at the base of the postabdominal claws, and the high heads.

Two of the discovered Cyclopoida species from the samples are new for the Albanian fauna - *Tropocyclops prasinus* and *Acanthocyclops vernalis*.

Because of the differences in the hydrological regime of the investigated waters the established species have various ecological characteristics. Some of them are typically pelagic forms like Rotifers *Euchlanis dilatata*, *Keratella cochlearis*, *K. quadrata*, Cladocerans *Sida cristalina*, *Bosmina longirostris* and Cyclopoida species - *Cyclops vicinus* found in the Skutari lake. Others are connected with small water bodies and with the bottom substrate. Such species are rotifer *Lecane (M.) arcuata*, cladocerans *Chydorus sphaericus*, *Alona gutata*, *Alona rectangula*, *Daphnia pulex pulex* and the Cyclopoid *Tropocyclops prasinus*. The greatest variety of species composition was established in the zooplankton from the Skutari lake - 19 species. The poorest samples were from a little well at Stani I Drugomiri place - 1590 m.a.s.l. containing only *Gammarus* sp. The rest of the samples contain between three and ten species.

Most of the established species have a cosmopolitan distribution. Only some of the species are somewhat stenobiotic as for example *Tropocyclops prasinus* found in the artificial lake in the Rasma village. This species is characteristic for the warm waters in the small lakes and ponds, during the summer season. It is monocyclic.

Three mature female individuals found in the lake N 3 are between 750 and 820 μm long. The lateral parts of the fifth toracal segment are covered by a lot of setae. The furcal rami are short and slightly divergent in their distal parts. They are approximately three times longer as broad. Of the middle apical seta of the furca, the inner one is five times longer than the furcal rami. The lateral furcal setae are well developed. They are attached to the back 1/3 part of the furcal rami. The antennulae are comparatively short, twelve membered, with dented hyaline membranes of the distal joint. They reach up to the third cephalotoracal segment. P5 is - one somited with three appendages a thin spine in the middle and two setae. The inner one apically situated and the external one subapically situated.

Another species is *Acanthocyclops vernalis* - characteristic for higher altitudes. It was found in the higher artificial ponds situated above Rasma village. In each of the samples in which this species was founded, the number of the individuals was comparatively large. In Bulgaria this species is common for the high mountain glacial lakes in Rila and Pirin mountains over 2095 m.a.s.l. (NAIDENOV, 1975).

The low temperature of the water and the unsustainable hydrological regime are the two main factors responsible for the poor fauna found in the spring and well at the Stani I Drugomiri place. Only four benthic taxa were found in them. The highest situated ponds and sources are much poorer in animals than the lower situated Skutari lake.

References

- BREM V., E. ZADERBAUER. 1905. Das Septemberplankton des Skutari-Sees. - Verh. K.k. zool.-bot. Ges., Wien, 9 (2): 55-150
- GESSNER F. 1934. Limnologische Untersuchungen am Skadar-(Skutari) See. - Glasnik botan. Zavodu I baste Univ. Beograd, 3 (11): 80-115.
- NAIDENOV W. 1975. Biologische Eigenheiten der glazialen Hydrofauna aus den Gebirgen Rila und Pirin (Bulgarien) - Symp. Biol. Hung., 15: 281-284.
- NEDELKOVIC R. 1959. Skadarsko jezero. - Monografija 4. Biol. inst. NRS, Beograd, 1-235
- PANDOURSKI I. 1997. Cyclopoides (Crustacea, Copepoda) des eaux souterraines de l'Albanie. - Acta zool. Bulg., 49: 64-67.
- PETKOVSKI T. 1961. Zur Kenntnis der Crustaceen des Skadar (Scutari)-Sees. - Acta Mus. Maced. Sci. Natur., 7: 29-52.
- VERESICHAGIN I. 1912. Cladocera from the lake Skutari (Monte Negro) - Works of the Zoological laboratory of HM Varshava Univ., 15: 56-98. (In Russian).

Received on 2.12.1998

Author's address:
Dimitar Kozuharov
Institute of Zoology
1, Tsar Osvoboditel Blvd
1000 Sofia, Bulgaria
e-mail: zoology@hgciat.acad.bg

Зоопланктонът от някои водни басейни в Северна Албания с различна големина и надморска височина

Димитър КОЖУХАРОВ

(Р е з ю м е)

Изследвани са зоопланктонни организми от осем различни по размер и хидрологичен режим водни басейна в Северна Албания. Обектите за изследване са разположени от 16 м н.в. до 1560 м н.в. От изследваните обекти са установени 14 таксона Rotatoria, 11- таксона Cladocera, 6 таксона Cuscleroidea (табл. 1). От намерените циклопоиди два вида не са съобщавани за Албания - *Tropocyclops prasinus* и *Acanthocyclops vernalis*, Клагоцерамата *Daphnia galeata*, намерена в изкуствено езеро в, също не е съобщавана за Албания. Най-богати са пробите от Шкодренското езеро, а най-малко видове, при това само бентосни, са установени в пробите от кладенчето в местността Стани и Другомири - 1560 м н.в.